

EGSIEM combination service: combination of GRACE monthly K-band solutions on normal equation level

U. Meyer¹, Y. Jean¹, A. Jäggi¹, D. Arnold¹
and the EGSIM-ACs^{2,3,4}

¹ Astronomical Institute, University of Bern

² Helmholtz Centre Potsdam, GFZ German Research Centre for Geosciences

³ Institute for Theoretical and Satellite Geodesy, TU Graz

⁴ Groupe de Recherche de Géodésie Spatiale, Toulouse

EGU General Assembly 2017

Vienna


April 24-28, 2017

Contents

- Motivation
- Individual Contributions
- Noise Assessment
- Combination on Normal Equation Level
- Conclusions


Motivation

Degree Amplitudes of Anomalies 01/2006: orders 0 - 29
SH coefficients – model fit of secular/seasonal variations




Motivation

Degree Amplitudes of Anomalies 01/2006: orders 0 - 29
SH coefficients – model fit of secular/seasonal variations




Motivation

Degree Amplitudes of Anomalies 01/2006: orders 0 - 29
SH coefficients – model fit of secular/seasonal variations




Motivation


Degree Amplitudes of Anomalies 01/2006: orders 0 - 29
SH coefficients – model fit of secular/seasonal variations




Individual Contributions




Individual Contributions



Individual Contributions



Individual Contributions



Individual Contributions

Why are formal errors so different?

Formal errors depend on the noise model applied!

**Error propagation of
kinematic orbits and
K-band observations**




Optimistic

**Realistic
(empirical)**


**Errors of observations:
GPS, K-band, accelerometers,
star cameras**



**Errors of background models
and de-aliasing: ocean tides,
short periodic atmosphere
and ocean variations (AOD)**




Noise Assessment



Anomalies: —
differences to model —



Noise Assessment




Anomalies: ———
differences to model

Differences: - - -
differences to mean


RMS of anomalies





RMS of differences to mean





Noise Assessment




Differences to mean
to derive relative
weights.  

Anomalies over quite
regions to indepently
assess quality.  


RMS of anomalies




RMS of differences to mean



Variance component estimation on solution level



Variance component estimation on solution level taking into account all SH coefficients up to degree and order 80 with equal weight.



RMS of anomalies restricted to ocean areas as quality criterion.

Combination on Normal Equation Level

Achieve equal impact of individual contributions on pairwise combinations:


$$(\mathbf{N}_{\text{ref}} + w_i \mathbf{N}_i) d\mathbf{x} = \mathbf{b}_{\text{ref}} + w_i \mathbf{b}_i$$

The impact is measured by:

$$\text{RMS}_i = \sqrt{\frac{\sum_{l,m} (K_{l,m}^{\text{comb}} - K_{l,m}^i)^2}{n_{\text{coef}}}}$$

Equal impact is achieved for: $\frac{\text{RMS}_i}{\text{RMS}_{\text{ref}}} = 1$


Combination on Normal Equation Level




equalizing weight

GRGS	1.60
GFZ	1.00
AIUB	7.81
ITSG	2.21

Combination: 2006/01




Combination: 2006/01



Solution:	weight
GRGS	0.14
GFZ	0.19
AIUB	0.29
ITSG	0.38

Combination: 2006/01




equalizing weight

GRGS	1.60
GFZ	1.00
AIUB	7.81
ITSG	2.21


Solution: weight

GRGS	0.14
GFZ	0.19
AIUB	0.29
ITSG	0.38

L3-Products: www.egsiem.eu -> Data -> EGSIEM-Plotter



L3-Products: www.egsiem.eu -> Data -> EGSiEM-Plotter



Conclusions

- EGSiem monthly gravity field combination on NEQ-level is operational.
- Noise assessment by variance component estimation on solution level.
- Relative weights based on noise levels.
- The EGSiem combination service provides two test years (2006 + 2007):
 - SH-coefficients (Level-2): www.icgem.de
 - grids and de-aliasing (Level-3): www.egsiem.eu